Intervention: Booster sessions

Finding: Sufficient evidence for effectiveness

Potential partners to undertake the intervention:	
Nonprofits or local coalitions	☐Businesses or labor organizations
Schools or universities	☐Media
Health care providers	Local public health departments
☐State public health departments	□Policymakers
☐ Hospitals, clinics or managed care organizations	Other:

Background on the intervention:

Booster sessions take place after an individual or group has completed the primary intervention's curriculum. Booster sessions provide an opportunity to reinforce and build on the messages and skills learned during the primary intervention.

Findings from the systematic reviews:

Systematic reviews provide sufficient evidence to support the use of booster sessions in alcohol and other drug abuse prevention programs.

Limitations/Comments:

The number of booster sessions that accompanied primary intervention curricula varied significantly among studies. Thus, the frequency and timing of booster sessions necessary to produce positive effects are unclear. In addition, the positive effects associated with booster programs may depend on other program characteristics.

References:

Cuijpers P. Effective ingredients of school-based drug prevention programs. A systematic review. Addict. Behav. 2002 Nov-Dec; 27(6):1009-1023.

McBride N. A systematic review of school drug education. Health Educ. Res. 2003 Dec; 18(6):729-742.

Midford R, Munro G, McBride N, Snow P, Ladzinski U. Principles that underpin effective school-based drug education. J. Drug Educ. 2002; 32(4):363-386.

Thomas H, Siracusa L, Ross G, Beath L, Hanna L, Michaud M, et al. Effectiveness of school-based interventions in reducing adolescent risk behaviour: A Systematic Review of Reviews. March 1999.